

Granular cell tumour of the appendix: an unexpected location

Joana Inês Martins Madeira, Rui Caetano Oliveira , Mário Rui Silva, Maria Augusta Cipriano

Pathology, Centro Hospitalar e Universitário de Coimbra EPE, Coimbra, Portugal

Correspondence to

Dr Joana Inês Martins Madeira; ji_madeira@hotmail.com

Accepted 29 November 2022

DESCRIPTION

Granular cell tumour (GCT) is an uncommon but usually benign lesion of neural/Schwann origin. This tumour type is most frequently found in skin, subcutaneous tissue and oral cavity, mostly in black middle-aged women.¹ This lesion type can involve any organ, including the gastrointestinal (GI) tract. The appendicular involvement though is extremely rare.

We report a case of a pregnant female patient in her early 30s who went to the emergency department after 2 days of enduring persistent pelvic pain. The blood tests indicated leucocytosis, neutrophilia and elevated C reactive protein concentration.

Abdominal ultrasound examination revealed an outer appendiceal diameter of 8 mm, with a markedly thickened and stratified wall. The increased density of the surrounding fat tissue was consistent with phlegmon.

A laparoscopic appendectomy was performed.

Careful gross examination showed that the appendix was increased in size, the appendiceal wall was markedly thickened and the lumen filled with faecal material.

The complete histological analysis of the specimen revealed a well-circumscribed and unencapsulated

nodule with 0.5 cm in diameter confined to the submucosa. There was also a florid granulomatous chronic inflammation measuring 1.2 cm extending to the tip of the appendix (as shown in [figure 1A](#)). This granulomatous inflammation was located in the wall and involved its entire thickness, except the overlying mucosa. The multiple, coalescent microgranulomas (non-necrotising granulomas) were composed only of epithelioid histiocytes. The authors want to emphasise that there was no clinical or imaging procedure of the thorax, which may constitute evidence for tuberculosis diagnosis. The surrounding tissues showed reactive and hyperplastic mucosa, with no atrophy. There were no acute appendicitis lesions.

The nodule was composed of nests of epithelioid cells with abundant granular eosinophilic cytoplasm and central small round nuclei (as shown in [figure 1C](#)). The tumour was devoid of cytological atypia and necrosis.

Immunohistochemistry revealed S-100 protein strong and diffuse positivity, suggestive of Schwann cell origin, and it was not expressed by the granulomatous epithelioid cells. The GCT cells also displayed SOX-10 nuclear expression (as shown in [figure 1D](#)) and CD68 membrane expression. CD68




Figure 1 (A) Submucosal GCT (arrow) was contiguous with a chronic granulomatous inflammation zone (star), (H&E, $\times 20$). (B) The tumour cells and granulomas without necrosis found in the appendix wall were immunoreactive for CD68. (C) The round to oval tumour cells displayed a granular cytoplasm (H&E, $\times 100$). (D) The neoplastic cells typically expressed nuclear SOX-10 ($\times 100$). GCT, granular cell tumour.



© BMJ Publishing Group Limited 2022. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Madeira JIM, Oliveira RC, Silva MR, et al. *BMJ Case Rep* 2022;**15**:e251058. doi:10.1136/bcr-2022-251058

Table 1 Clinical characteristics of patients reported in the literature with GCT of the appendix

Case number	Author(s), year	Age, gender	Clinical symptoms	Radiology assumption	Size of tumour (cm)
1	Wanick, 1956 ⁴	30s, female	Acute pain in right iliac fossa, vomiting and diarrhoea	Acute appendicitis	0.5 and 1 (double nodules)
2	Hausman, 1963 ⁵	40s, male	NP	NP	0.8
3	Apisarnthanarax, 1981 ⁶	30s, female	NP	NP	5
4	Sarma <i>et al</i> , 1984 ⁷	50s, male	Asymptomatic	NP	0.5
5	Fried <i>et al</i> , 1984 ⁸	30s, male	NP	NP	0.8
6	Pipeleers-Marichal <i>et al</i> , 1990 ⁹	40s, male	Asymptomatic	NP	4
7	Kaltschmidt <i>et al</i> , 1992 ¹⁰	30s, male	NP	Acute appendicitis	NP
8	Gavelli <i>et al</i> , 2005 ¹¹	40s, male	Acute pain in right iliac fossa and fever	Acute appendicitis	0.5
9	Moreno Gijon <i>et al</i> , 2009 ¹²	30s, female	Acute pain in right iliac fossa	Endometrioma	NP
10	Saleh <i>et al</i> , 2009 ¹³	60s, female	Asymptomatic	NP	0.5
11	Singhi and Montgomery 2010 ¹⁴	40s, female	NP	Acute appendicitis	0.6
12	Zoccali, 2011 ¹⁵	late adolescence, male	Acute pain in right lower quadrant	Abscess	3.5
13	Roncati <i>et al</i> , 2013 ¹	40s, female	Acute pain in right lower quadrant, pain at defecation, tenesmus, rectal bleeding and haematuria	Phlegmon	0.2
14	Roy and Goswami, 2015 ¹⁶	20s, male	Acute pain in right iliac fossa and vomiting	NP	NP
15	Allison and Rao, 2020 ²	50s, female	NP	NP	NP
16	Current case	30s, female	Acute pelvic pain	Phlegmon	0.5

Source: PubMed.
GCT, granular cell tumour; NP, not provided.

pointed out to the granulomas found in the appendix wall (as shown in [figure 1B](#)).

The postoperative period was uneventful, and the patient was discharged after 1 week.

Less than 15% of GCTs occur in the GI tract, with only 15 cases being reported in the appendix so far.^{1 2}

A detailed analysis ([table 1](#)) of the features of GCTs in the appendix reveals that the male/female ratio was 1:1, the mean age was that of 41 years old and the average size of the lesions was 1.4 cm. The patients' clinical characteristics, including from the patient reported in this article, can be consulted in [table 1](#).

The pathogenesis of GCTs remains unclear; however, the medical literature data suggest that a chronic inflammation, surrounding the GCT, is an antecedent condition that may favour its appearance.

In fact, chronic and unresolved inflammation promotes immunosuppression and so it has been associated with an increased risk of developing both benign and malignant tumours.

Furthermore, published reports indicate that GCT in the appendix is a lesion which reflects local reactive changes of the neural/Schwann cells rather than a true neoplasm.³

This case documents a GCT probably arising from a granulomatous appendicitis background, which is a rare entity. This chronic inflammation corresponds to less than 2% of appendectomies as a cause of appendicitis.

Surgical intervention with wide local excision and complete resection is still the best treatment for GCTs in the GI tract, with excellent outcomes for benign lesions and with a low rate of recurrence or malignant transformation.

Learning points

- Granular cell tumour (GCT) is an uncommon benign lesion of neural/Schwann origin.
- The appendicular involvement is extremely rare.
- We report a case of granulomatous appendicitis and a GCT as an incidental finding, whose symptoms and analytical parameters mimic an appendicitis.

There are no data in the literature that correlate with the emergence of GCT in the lower GI tract and the risk of developing colorectal cancer.

Previous reports suggest that regardless of the initial nature of the lesion, all patients should be followed to prevent recurrence and distant metastasis.

Contributors JIMM and RCO—drafting and revision of preliminary and final versions, and approval of the final version. MRS and MAC—histology, revision of preliminary and final versions, and approval of the final version.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

Case reports provide a valuable learning resource for the scientific community and can indicate areas of interest for future research. They should not be used in isolation to guide treatment choices or public health policy.

ORCID iD

Rui Caetano Oliveira <http://orcid.org/0000-0002-7202-8059>

REFERENCES

- 1 Roncati L, Manco G, Italia S, *et al*. Granular cell tumor of the appendix: a new case and review of the literature. *Springerplus* 2013;2:649.
- 2 Allison D, Rao D. Granular cell tumor confined to the muscularis propria of the appendix: a case report. *Am J Clin Pathol* 2020;154:S57–8.
- 3 Vered M, Carpenter WM, Buchner A. Granular cell tumor of the oral cavity: updated immunohistochemical profile. *J Oral Pathol Med* 2009;38:150–9.
- 4 Wanick LR. [Histogenesis of granular cell myoblastoma of the appendix; case presentation]. *Mem Inst Oswaldo Cruz* 1956;54:125–34. doi:10.1590/s0074-02761956000100007
- 5 HausmanR. Granular cells in musculature of the appendix. *Arch Pathol* 1963;75:360–72.
- 6 Apisarnthanarax P. Granular cell tumor. An analysis of 16 cases and review of the literature. *J Am Acad Dermatol* 1981;5:171–82. doi:10.1016/s0190-9622(81)70085-9
- 7 Sarma DP, Simmons GT, Rodriguez FH, *et al*. Granular cell tumor of the appendix. *J Surg Oncol* 1984;27:219–21. doi:10.1002/jso.2930270403
- 8 Fried KS, Arden JL, Gouge TH, *et al*. Multifocal granular cell tumors of the gastrointestinal tract. *Am J Gastroenterol* 1984;79:751–6.

- 9 Pipeleers-Marichal M, Goossens A, De Waele B, *et al.* Granular cell tumour of the appendix in a patient irradiated for a rectal carcinoma. *Virchows Arch A Pathol Anat Histopathol* 1990;417:177–80. doi:10.1007/BF02190537
- 10 Kaltschmidt A, Wershigora JW, Schuh D. [Granular cell tumor of the vermiform appendix. Case presentation with discussion of histogenesis]. *Zentralbl Pathol* 1992;138:55–64.
- 11 Gavelli A, Clément N, Marmoreale A, *et al.* [Granular cell tumor involving the appendix]. *Gastroenterol Clin Biol* 2005;29:211–3. doi:10.1016/s0399-8320(05)80744-4
- 12 Moreno Gijón M, Granero Castro P, Fernández García MS, *et al.* [Granular cell tumor of the appendix]. *Rev Esp Enferm Dig* 2009;101:512–5. doi:10.4321/s1130-01082009000700015
- 13 Saleh H, El-Fakharany M, Frankle M. Multiple synchronous granular cell tumors involving the colon, appendix and mesentery: a case report and review of the literature. *J Gastrointestin Liver Dis* 2009;18:475–83.
- 14 Singhi AD, Montgomery EA. Colorectal granular cell tumor: a clinicopathologic study of 26 cases. *Am J Surg Pathol* 2010;34:1186–92. doi:10.1097/PAS.0b013e3181e5af9d
- 15 Zoccali M, Cipriani N, Fichera A, *et al.* Acute appendicitis secondary to a granular cell tumor of the appendix in a 19-year-old male. *J Gastrointest Surg* 2011;15:1482–5. doi:10.1007/s11605-011-1536-4
- 16 Roy A, Goswami PK. Granular cell tumour of appendix – a case report. *Paripep-indian Journal Of Research* 2015;4.

Copyright 2022 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <https://www.bmj.com/company/products-services/rights-and-licensing/permissions/>
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

Customer Service

If you have any further queries about your subscription, please contact our customer services team on +44 (0) 207111 1105 or via email at support@bmj.com.

Visit casereports.bmj.com for more articles like this and to become a Fellow